RESEARCH BRIEFS

Short Reports on Current Research in Agricultural Sciences

CITRUS NEMATODE BIOTYPES


PRELIMINARY DATA on the distribution and prevalence of different biotypes of the citrus nematode in California show a scattered, wide distribution pattern for four biotypes. Biotypes are important in the selection and development of citrus nematode-resistant rootstocks. Some selections of trifoliate orange and certain citranges differ in their susceptibility to different biotypes of the citrus nematode. This survey is being continued.—R. C. Baines, Dept. of Nematology, U.C. River-

GRAPEFRUIT RIND BREAKDOWN

WITH INCREASED plantings of summer grapefruit, the problem of rind breakdown gains importance. Rind staining and/or concentric rind stipple are fairly common in certain years. However, during the 1973 season, a number of lots of fruit had a rind breakdown and/or decay either at packing or during shipment. A large committee of UC and Sunkist researchers, packinghouse managers, and a local laboratory manager has been assembled to make a long-term study of the problem. It appears from preliminary leaf analysis data that some groves with high nitrogen experienced more rind breakdown.—R. M. Burns, Cooperative Extension, Ventura County.

PESTICIDE

FARM WORKER SAFETY DATA







CALIFORNIA AGRICULTURE

Progress Reports of Agricultural Research, published monthly by the University of California Division of Agricultural Sciences.

Articles published herein may be republished or reprinted provided no advertisement for a commercial product is implied or imprinted.

address

141



A continuing program of research in many aspects of agriculture is carried on at University campuses, field stations, leased areas, and many temporary plots loaned by cooperating landowners throughout the state. Listed below are some of the projects currently under way, but on which no formal progress reports can yet be made.

WEED-EATING INSECTS

BIOLOGICAL CONTROL specialists at Berkeley are attempting to find otherwise harmless insect predators that will help to control Scotch broom, tansy ragwort, gorse, and hoary cress.

ARE CO-OPS EFFECTIVE?

BY TRACING the history of the major agricultural cooperatives in California, economists at Berkeley hope to provide an improved basis on which California farmers many use such associations as effective tools for solving marketing problems.

DECONTAMINATION OF FEEDSTUFFS

SCIENTISTS AT DAVIS are investigating pesticide residues and searching for ways to remove them or render them harmless by chemical or physical means, or by a combination of both. Research will be concerned primarily with the pesticides most likely to present problems in the more important feedstuffs and associated foods.